




## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>6</sup>:</b> <b>A61K 31/435, C07D 471/04</b>	<b>A1</b>	<b>(11) International Publication Number:</b> <b>WO 97/07800</b> <b>(43) International Publication Date:</b> 6 March 1997 (06.03.97)
<b>(21) International Application Number:</b> PCT/IB96/00756 <b>(22) International Filing Date:</b> 29 July 1996 (29.07.96) <b>(30) Priority Data:</b> 60/002,975                      29 August 1995 (29.08.95)                      US <b>(71) Applicant (for all designated States except US):</b> PFIZER INC. [US/US]; 235 East 42nd Street, New York, NY 10017 (US). <b>(72) Inventors; and</b> <b>(75) Inventors/Applicants (for US only):</b> ALLEN, Douglas, John, Meldrum [US/US]; 549 Ocean Avenue, New London, CT 06320 (US). JOSEPH, David, Bruning [US/US]; Unit 2N, 70 Farmington Avenue, New London, CT 06320 (US). NORRIS, Timothy [GB/US]; 27 Friar Tuck Drive, Gales Ferry, CT 06335 (US). <b>(74) Agents:</b> SPIEGEL, Allen, J. et al.; Pfizer Inc., 235 East 42nd Street, New York, NY 10017 (US).		<b>(81) Designated States:</b> AU, BG, BR, BY, CA, CN, CZ, HU, IL, IS, JP, KR, KZ, LK, LV, MX, NO, NZ, PL, RO, RU, SG, SI, SK, TR, UA, US, UZ, VN, European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).  <b>Published</b> <i>With international search report.</i>
<b>(54) Title:</b> ZWITTERIONIC FORMS OF TROVAFLOXACIN <b>(57) Abstract</b> <p>A zwitterionic form of trovafloxacin having formula (I) selected from the group consisting of its crystalline hygroscopic and non-hygroscopic polymorphs and pentahydrate and methods for their preparation. The invention further relates to methods of using, and pharmaceutical compositions comprising, the compounds of the invention for treatment of bacterial infections in mammals.</p> <div style="text-align: center;">  <p>(I)</p> </div>		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore
CH	Switzerland	LI	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SZ	Swaziland
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Latvia	TG	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	UG	Uganda
FI	Finland	MN	Mongolia	US	United States of America
FR	France	MR	Mauritania	UZ	Uzbekistan
GA	Gabon			VN	Viet Nam

5

ZWITTERIONIC FORMS OF TROVAFLOXACINBackground of the Invention


This invention relates to the naphthyridone antibiotic trovafloxacin. More particularly, it relates to polymorphs and the pentahydrate of the zwitterionic form of thereof having the formula I, below, and methods for their preparation. The invention further relates to methods of using, and pharmaceutical compositions comprising, the compounds of the invention for treatment of bacterial infections in mammals.

The antibacterial activity of the aforementioned naphthyridone antibiotic is described in United States Patent No. 5,164,402 [the '402 patent] and 5,229,396 issued 11/17/92 and 7/20/93, respectively, the disclosures of which are hereby incorporated herein by reference in their entirety. The foregoing patents are assigned in common with the present application.

The zwitterionic forms of trovafloxacin are useful for the administration of the drug as a suspension.

Summary of the Invention

According to a first embodiment of the invention there is provided a trovafloxacin zwitterionic crystal form having the formula



25

30

which is selected from the group consisting of

- a) a non hygroscopic first polymorph PI exhibiting the following characteristic X-ray powder diffraction pattern

35

-2-

5	<u>Peak no.</u>	<u>1</u>	<u>2</u>	<u>3</u>	<u>4</u>	<u>5</u>	<u>6</u>	<u>7</u>	<u>8</u>	<u>9</u>
	2_θ_(°) Cu	6.9	9.8	11.3	12.0	13.9	16.1	16.6	17.1	17.4
	d space	12.7	9.0	7.9	7.4	6.4	5.5	5.4	5.2	5.1
	<u>Peak no.</u>		<u>10</u>	<u>11</u>	<u>12</u>	<u>13</u>	<u>14</u>	<u>15</u>	<u>16</u>	<u>17</u>
	2_θ_(°) Cu		19.7	22.9	23.6	24.9	25.4	25.9	27.7	29.5
	d space		4.5	3.9	3.8	3.6	3.5	3.4	3.2	3.0

- 10 b) a hygroscopic second polymorph PII exhibiting the characteristic X-ray powder diffraction pattern

<u>Peak no.</u>	<u>1</u>	<u>2</u>	<u>3</u>	<u>4</u>	<u>5</u>	<u>6</u>	<u>7</u>	<u>8</u>
2_θ_(°) Cu	8.4	9.5	10.2	14.7	16.8	17.9	22.6	26.1
d space	10.6	9.3	8.7	6.0	5.3	5.0	3.9	3.4

15 and

- c) a pentahydrate, trovafloxacin zwitterion pentahydrate, exhibiting the characteristic X-ray powder diffraction pattern

20	<u>Peak no.</u>	<u>1</u>	<u>2</u>	<u>3</u>	<u>4</u>	<u>5</u>	<u>6</u>	<u>7</u>	<u>8</u>	<u>9</u>
	2_θ_(°) Cu	6.6	8.6	12.7	13.3	15.9	18.6	19.2	20.1	21.0
	d space	13.3	10.3	7.0	6.6	5.5	4.8	4.6	4.4	4.2
25	<u>Peak no.</u>	<u>10</u>	<u>11</u>	<u>12</u>	<u>13</u>	<u>14</u>	<u>15</u>	<u>16</u>	<u>17</u>	
	2_θ_(°) Cu	22.5	22.9	23.6	24.9	25.4	25.9	27.7	29.5	
	d space	4.0	3.9	3.8	3.6	3.5	3.4	3.2	3.0	

-3-

A second embodiment of the invention relates to a process for preparing a zwitterion, of trovafloxacin, of the formula I which is selected from the group consisting of a non hygroscopic polymorph PI, a hygroscopic polymorph PII and a pentahydrate thereof, as described above, comprising

5           A.     the steps of treating an aqueous suspension of a metastable form of the compound of the formula I

                  1)     with a nonpolar solvent followed by azeotropic removal of residual water and vacuum drying to form said hygroscopic polymorph PII which exhibits the characteristic X-ray powder diffraction pattern described in claim 1;

10               2)     with a polar solvent followed by azeotropic removal of residual water and vacuum drying; or

                  3)     with water and air drying the residue at an elevated temperature, removing the mother liquor and air drying the residue at room temperature to constant weight to form the pentahydrate; or

15           B)     treating the hygroscopic second polymorph PII with a refluxing polar solvent to form the non-hygroscopic first polymorph PI.

According to a third embodiment of the invention there is provided a process for preparing the metastable form of the zwitterion, of trovafloxacin, of the formula I, by


20           a)     treating an acid salt of trovafloxacin with a base to raise the pH of the mixture to between 7.5 and 8.5 at an elevated temperature, removal of the mother liquor, washing the crystals with water and drying the crystals under vacuum at about 35 to about 40 °C; or

                  b)     treating a compound of the formula  
25

30

-4-

5



II

10

6

wherein A is hydrogen or an amine protecting group such as *t*-butoxycarbonyl, benzyloxycarbonyl, (C<sub>1</sub>-C<sub>6</sub>)alkylcarbonyl and benzyl; and

B is hydrogen or a carboxylic acid protecting group selected from benzyl, *t*-butyl and (C<sub>1</sub>-C<sub>6</sub>) alkyl; with an amine and/or carboxylic acid deprotecting agent, respectively.

A fourth embodiment of the invention provides a method of treating bacterial infections in a mammal which comprises administering to said mammal a bacterial infection treating effective amount of a compound of formula I as described above.


According to a fifth embodiment of the invention there is provided a composition for treating bacterial infections in a mammal which comprises a bacterial infection treating effective amount of a compound of formula I and a pharmaceutically acceptable carrier.

#### Detailed Description of the Invention

The present invention relates to a compound comprising a stable zwitterionic form of the antibiotic trovafloxacin of the formula

30

-5-




More particularly, it is related to a compound of the formula I which is selected from

- a) a non hygroscopic first polymorph PI exhibiting the characteristic X-ray powder diffraction pattern described above;
- 15 b) a hygroscopic second polymorph PII exhibiting the characteristic X-ray powder diffraction pattern described above;
- and c) a pentahydrate, trovafloxacin zwitterion pentahydrate, exhibiting the characteristic X-ray powder diffraction pattern described above.

The invention also relates to processes for the preparation of the compounds

20 of the formula I as illustrated in the following schemes.


-6-

SCHEME 1



-7-

## SCHEME 2




metastable forms of zwitterion

-8-

As shown in Scheme 1 a trovafloxacin salt 1, wherein X is an anion selected from those formed from mineral acids such as hydrochloric, sulfuric, nitric and phosphoric; organic acids such as sulfonic acids, e. g. benzenesulfonic (besylic), p-toluenesulfonic (PTSA, tosylic), methanesulfonic (MSA, mesylic) and trifluoromethanesulfonic (triflic); and carboxylic acids e.g., acetic, propionic, benzoic, citric, tartaric, maleic, fumaric, succinic and malic, is converted to a metastable zwitterionic form 2 by raising the pH of a slurry comprising compound 1 to a pH of between about 7.5 and 8.5 at a temperature in the range of about 45 to about 55°C using an aqueous basic solution. A preferred salt is the mesylate. The bases useful in the practice of this aspect of the invention include inorganic bases such as alkali or alkaline earth hydroxides, carbonates and bicarbonates and organic bases such as tri(C<sub>1</sub>-C<sub>6</sub>)alkyl amines, pyridine and morpholine. A preferred aqueous base is saturated sodium bicarbonate. The wet product is then dried to constant weight, *in vacuo*, at a temperature from about 35 to about 40°C.

Alternatively, as shown in scheme 2, compound 2 may be prepared directly from protected precursors 6, of the trovafloxacin salts 1, of the formula



wherein A is hydrogen or an amine protecting group such as *t*-butoxycarbonyl, benzyloxycarbonyl, (C<sub>1</sub>-C<sub>6</sub>)alkylcarbonyl and benzyl; and

B is hydrogen or a carboxylic acid protecting group selected from benzyl, *t*-butyl and (C<sub>1</sub>-C<sub>6</sub>) alkyl; with an amine and/or carboxylic acid deprotecting agent, respectively.

-9-

A preferred compound 6, wherein A is hydrogen and B is ethyl, is converted to compound 2 by treatment with a solution of NaOH in a polar solvent at an elevated temperature. A preferred solvent is methanol and the temperature is the reflux temperature of the solvent. The pH of the solution was then adjusted to between about  
5 6.5 and 8.0 with dilute HCl and saturated aqueous NaHCO<sub>3</sub> was then added to adjust the pH to between about 7.5 and 8.5. The product was recovered as indicated above.

Metastable trovafloxacin zwitterion 2 is converted to hygroscopic polymorph PII, 4, by treatment with a non polar solvent such as a hydrocarbon. A preferred hydrocarbon is hexanes. Residual water is removed azeotropically and the product  
10 dried at about 35 to about 40°C under vacuum. Solvents useful for the azeotropic removal of water traces include non-polar aliphatic hydrocarbons, such as hexanes and octanes, and aromatic hydrocarbons such as benzene and toluene. Preferred solvents are the aliphatic hydrocarbons, most preferably hexanes.

Non hygroscopic polymorph PI 3, can be prepared from compound 2 by  
15 treatment with a polar solvent followed by azeotropic removal of water and vacuum drying at about 30 to about 40°C. Polar solvents useful for this conversion include (C<sub>1</sub>-C<sub>6</sub>)alkyl esters of (C<sub>2</sub>-C<sub>6</sub>)alkylcarboxylic acids and (C<sub>1</sub>-C<sub>6</sub>)alkanols. A preferred solvent is ethyl acetate.

Alternatively, compound 3 can be prepared from compound 4 by treating  
20 compound 4 with a refluxing polar solvent, as described above. A preferred solvent is ethyl acetate.

Compound 5, the pentahydrate of the compound of formula I, is prepared by air drying the wet crystals of compound 1, at room temperature, until constant weight is achieved. Alternatively, compound 5 may be prepared from compound 4 by  
25 treatment with water until a constant water uptake has been obtained. Compound 3 is not converted to compound 5 by exposure to water.

The antibacterial compounds of the invention, i.e., polymorph PI, polymorph PII and the pentahydrate (hereafter "the active compounds") are useful in the treatment of animals and humans having a broad spectrum of bacterial infections. They are  
30 particularly useful in treating gram-positive bacterial strains.

The active compounds may be administered alone, but will generally be administered in a mixture with a pharmaceutical carrier selected with regard to the intended route of administration and standard pharmaceutical practice. For example, they can be administered orally or in the form of tablets containing such excipients as

-10-

starch or lactose, or in capsules either alone or in admixture with excipients, or in the form of elixirs or suspensions containing flavoring or coloring agents. In the case of animals, they are advantageously contained in an animal feed or drinking water in a concentration of about 5 to about 5000 ppm, preferably about 25 to about 500 ppm.

- 5 They can be injected parenterally, for example, intramuscularly, intravenously or subcutaneously. For parenteral administration, they are best used in the form of a sterile aqueous solution which can contain other solutes, for example, enough salt or glucose to make the solution isotonic. In the case of animals, the compounds of formula I can be administered intramuscularly or subcutaneously at dosage levels of  
10 about 0.1 to about 50 mg/kg/day, advantageously about 0.2 to about 10 mg/kg/day given in a single daily dose or up to 3 divided doses.

- The active compounds can be administered to humans, for the treatment of bacterial diseases by either oral or parenteral routes. They may be administered orally at dosage levels of about 0.1 to 500 mg/kg/day, advantageously 0.5-50 mg/kg/day  
15 given in a single dosage or up to 3 divided dosages. For intramuscular or intravenous administration, dosage levels are about 0.1-200 mg/kg/day, advantageously 0.5-50 mg/kg/day. While intramuscular administration may be a single dose or up to 3 divided doses, intravenous administration can include a continuous drip. Variations will necessarily occur depending on the weight and condition of the subject being treated  
20 and the particular route of administration chosen as will be known to those skilled in the art.

- The active compounds may be administered alone, but will generally be administered in a mixture with a pharmaceutical carrier selected with regard to the intended route of administration and standard pharmaceutical practice. For example,  
25 they can be administered orally or in the form of tablets containing such excipients as starch or lactose, or in capsules either alone or in admixture with excipients, or in the form of elixirs or suspensions containing flavoring or coloring agents. In the case of animals, they are advantageously contained in an animal feed or drinking water in a concentration of about 5 to about 5000 ppm, preferably about 25 to about 500 ppm.  
30 They can be injected parenterally, for example, intramuscularly, intravenously or subcutaneously. For parenteral administration, they are best used in the form of a sterile aqueous solution which can contain other solutes, for example, enough salt or glucose to make the solution isotonic. In the case of animals, the compounds of formula I can be administered intramuscularly or subcutaneously at dosage levels of

-11-

about 0.1 to about 50 mg/kg/day, advantageously about 0.2 to about 10 mg/kg/day given in a single daily dose or up to 3 divided doses. The active compounds can be administered to humans by either oral or parenteral routes, and may be administered orally at dosage levels of about 0.1 to 500 mg/kg/day, advantageously 0.5-50 mg/kg/day given in a single dosage or up to 3 divided dosages. For intramuscular or intravenous administration, dosage levels are about 0.1-200 mg/kg/day, advantageously 0.5-50 mg/kg/day. While intramuscular administration may be a single dose or up to 3 divided doses, intravenous administration can include a continuous drip. Variations will necessarily occur depending on the weight and condition of the subject being treated and the particular route of administration chosen as will be known to those skilled in the art.

The antibacterial activity of the compounds of the invention is shown by testing according to the Steer's replicator technique which is a standard *in vitro* bacterial testing method described by E. Steers et al., Antibiotics and Chemotherapy, 9, 307 (1959).

The following examples illustrate the methods and compounds of the present invention. It will be understood, however, that the invention is not limited to the specific examples.

#### Example 1

##### Trovafloxacin zwitterion, metastable form

A. Trovafloxacin mesylate (prepared according to Example 13B of the '402 patent) (20 g) was stirred with demineralized water (100 mL). The crystal slurry was heated to about 50 °C and the slurry adjusted to a pH of about 8.0 by addition of saturated sodium bicarbonate solution. The slurry was held at about 50 °C for 30 minutes, allowed to cool to about 25 °C and stirred at this temperature for 30 minutes. The crystals were isolated by filtration and washed with demineralized water (27 mL). The wet crystals were suspended in demineralized water (100 mL) and stirred for about 1 hour at about 50 °C, then cooled to about 20°C and stirred at this temperature for about 1 hour. The crystals were filtered from the mother liquor, washed with demineralized water (about 27 mL) and dried to constant weight under vacuum at about 40 °C to yield the title product which contained 2.5 % residual water by analysis. Yield 16.25 g, 97 %.

B. The ethyl ester of trovafloxacin (prepared according to the method of co-pending United States patent application serial number 08/490827, filed June 15, 1995,

-12-

the disclosures of which are hereby incorporated herein by reference in its entirety. The foregoing application is assigned in common with the present application.

(10 g) was stirred with methanol (75 mL), water (25 mL) and sodium hydroxide pellets (1.8 g). The resultant mixture was heated to reflux at about 72 °C to form a solution.

- 5 The solution was cooled to about 25 °C and the pH adjusted to about 7.5, by addition of 6N hydrochloric acid, to form a slurry. Saturated sodium bicarbonate solution (50 mL) was added and the slurry stirred for 30 minutes at about 25 °C. The title product was isolated and washed with water (20 mL) and dried under vacuum about 45 °C. Yield 7.72 g, 82.5 %.

10

### Example 2

#### Trovafloxacin zwitterion polymorph PI (non hygroscopic form)

- Trovafloxacin mesylate, (75 g) was stirred with demineralised water (375 mL). The crystal slurry was heated to about 50 °C and the slurry adjusted to a pH of about 8.0 by addition of saturated sodium bicarbonate solution. The slurry was held at about
- 15 50 °C for 30 minutes, allowed to cool to about 25 °C and stirred at this temperature for 30 minutes. Crystals were isolated by filtration and washed with demineralised water (100 mL). The wet crystals were suspended in demineralised water (375 mL) and stirred for 1 hour at about 50 °C, then cooled to about 20 °C and stirred at this temperature for about 1 hour. The crystalline product was filtered from the mother
- 20 liquor and washed with demineralised water (about 100 mL). The wet crystals were stirred with ethyl acetate (1125 mL) and the resultant slurry heated to reflux and the water azeotropically removed. The essentially anhydrous slurry was cooled to about 25 °C, the crystals were isolated by filtration and dried under vacuum at 40 °C until all the solvent had been removed to provide the title product. Yield 60.9 g, 94 %.
- 25 The product is characterized by the X-ray powder diffraction pattern described above.

### Example 3

#### Trovafloxacin zwitterion hygroscopic polymorph PII

- The title product of Example 1, paragraph A, (5 g) was mixed with hexanes (150
- 30 mL) to form a slurry. The slurry was heated to reflux and traces of residual water were removed azeotropically. After 4 hours at reflux the crystal slurry was cooled to about 25 °C, isolated by filtration and dried to constant weight under vacuum at about 40 °C. Yield 4.7 g, 94 %. The title product was characterized by the X-ray powder diffraction pattern described above.

-13-

Example 4Trovafloxacin zwitterion pentahydrate


Trovafloxacin mesylate (50 g) was stirred with demineralised water (250 mL). The crystal slurry was heated to 50 °C and the slurry adjusted to a pH of about 8.0 by addition of saturated sodium bicarbonate solution. The slurry was held at about 50 °C for 30 minutes, allowed to cool to about 25 °C and stirred at this temperature for 30 minutes. The crystals were isolated by filtration and washed with demineralised water (70 mL). The wet crystals were suspended in demineralised water (250 mL) and stirred for 1 hour at about 50 °C, then cooled to about 20°C and stirred at this temperature for about 1 hour. The crystalline product was filtered from the mother liquor, washed with demineralised water (about 70 mL). The wet crystals were air dried to constant weight at room temperature to yield the title product which contained 17.6 % water by analysis. Yield 48.4 g, 84 %

The title product was characterized by the X-ray powder diffraction pattern described above.

-14-

CLAIMS

1. A trovafloxacin zwitterionic crystal form having the formula



selected from the group consisting of

- 15 a) a non hygroscopic first polymorph PI exhibiting the characteristic X-ray powder diffraction pattern

20

Peak no.	1	2	3	4	5	6	7	8	9
2 $\theta$ (°) Cu	6.9	9.8	11.3	12.0	13.9	16.1	16.6	17.1	17.4
d space	12.7	9.0	7.9	7.4	6.4	5.5	5.4	5.2	5.1
Peak no.	10	11	12	13	14	15			
2 $\theta$ (°) Cu	19.7	20.3	21.2	22.8	23.8	26.3			
d space	4.5	4.4	4.2	3.9	3.7	3.4			

- 25 b) a hygroscopic second polymorph PII exhibiting the characteristic X-ray powder diffraction pattern

30

Peak no.	1	2	3	4	5	6	7	8
2 $\theta$ (°) Cu	8.4	9.5	10.2	14.7	16.8	17.9	22.6	26.1
d space	10.6	9.3	8.7	6.0	5.3	5.0	3.9	3.4



-15-

and

c) a pentahydrate, trovafloxacin zwitterion pentahydrate, exhibiting the characteristic X-ray powder diffraction pattern

5	<u>Peak no.</u>	<u>1</u>	<u>2</u>	<u>3</u>	<u>4</u>	<u>5</u>	<u>6</u>	<u>7</u>	<u>8</u>	<u>9</u>
	2_θ_(°) Cu	6.6	8.6	12.7	13.3	15.9	18.6	19.2	20.1	21.0
	d space	13.3	10.3	7.0	6.6	5.5	4.8	4.6	4.4	4.2
10	<u>Peak no.</u>		<u>10</u>	<u>11</u>	<u>12</u>	<u>13</u>	<u>14</u>	<u>15</u>	<u>16</u>	<u>17</u>
	2_θ_(°) Cu		22.6	22.9	23.6	24.9	25.4	25.9	27.7	29.5
	d space			3.9	3.8	3.6	3.5	3.4	3.2	3.0

2. The compound according to claim 1 consisting of said non hygroscopic first polymorph PI.

15 3. The compound according to claim 1 consisting of said hygroscopic second polymorph PII.

4. The compound according to claim 1 consisting of said pentahydrate.

5. The compound according to claim 1 consisting of said metastable form.


6. A process for preparing a compound of the formula

20

-16-

5

10



selected from the group consisting of a non hygroscopic polymorph PI, a hygroscopic polymorph PII and a pentahydrate thereof comprising

A. the steps of treating an aqueous suspension of a metastable form of the compound of the formula I

1) with a nonpolar solvent followed by azeotropic removal of residual water and vacuum drying to form said hygroscopic polymorph PII which exhibits the characteristic X-ray powder diffraction pattern described in claim 1;

2) with a polar solvent followed by azeotropic removal of residual water and vacuum drying; or

3) with water and air drying the residue at an elevated temperature, removing the mother liquor and air drying the residue at room temperature to constant weight to form the pentahydrate; or


B. by treating the hygroscopic second polymorph PII with a refluxing polar solvent to form the non-hygroscopic first polymorph PI.

7. The process of claim 6 wherein the metastable form, of the compound of formula I, is prepared by

a) treating an acid salt of trovafloxacin with a base to raise the pH of the mixture to between 7.5 and 8.5 at an elevated temperature; or

b) by treating a compound of the formula

-17-



wherein A is hydrogen or an amine protecting group such as *t*-butoxycarbonyl, benzyloxycarbonyl, (C<sub>1</sub>-C<sub>6</sub>)alkylcarbonyl and benzyl; and B is hydrogen or a carboxylic acid protecting group selected from benzyl, *t*-butyl and (C<sub>1</sub>-C<sub>6</sub>) alkyl; with an amine and/or carboxylic acid deprotecting agent, respectively.

8. The process of claim 6 step a) wherein the non solvent is hexanes.
9. The process of claim 6 step b) wherein the polar solvent is ethyl acetate.
- 20 10. A method for treating bacterial infection in a mammal which comprises administering to said mammal a bacterial infection treating effective amount of the compound of claim 1.
11. The method of claim 10 wherein said compound is non hygroscopic first polymorph PI.
- 25 12. The method of claim 10 wherein said compound is hygroscopic second polymorph PII.
13. The method of claim 10 wherein said compound is trovafloxacin zwitterion pentahydrate.
14. A composition for treating bacterial infections in a mammal which comprises a bacterial infection treating effective amount of a compound of claim 1 and a pharmaceutically acceptable carrier.
- 30 15. The composition of claim 14 wherein said compound is the non hygroscopic first polymorph PI.

-18-

16. The composition of claim 14 wherein said compound is the hygroscopic second polymorph PII.

17. The composition of claim 14 wherein said compound is trovafloxacin zwitterion pentahydrate.

5 18. The composition of claim 14 wherein said composition is a suspension.

19. The method of claim 7 wherein A is hydrogen or an amine protecting group such as *t*-butoxycarbonyl, benzyloxycarbonyl, (C<sub>1</sub>-C<sub>6</sub>)alkylcarbonyl and benzyl.

20. The method of claim 7 wherein said compound is B is hydrogen and A is selected from benzyl, *t*-butyl and (C<sub>1</sub>-C<sub>6</sub>)alkyl.

## INTERNATIONAL SEARCH REPORT

Int. .onal Application No  
PCT/IB 96/00756A. CLASSIFICATION OF SUBJECT MATTER  
IPC 6 A61K31/435 C07D471/04

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C07D A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	US,A,5 229 396 (PFIZER INC.) 20 July 1993 cited in the application see examples 12,13 ---	1
A	US,A,5 164 402 (PFIZER INC.) 17 November 1992 cited in the application see examples 12,13 -----	1

☐ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

## \* Special categories of cited documents :

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "B" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

10 September 1996

Date of mailing of the international search report

20.09.96

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+ 31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+ 31-70) 340-3016

Authorized officer

Van Bijlen, H

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/IB 96/00756

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US-A-5229396	20-07-93	WO-A- 9102526	07-03-91
		US-A- 5164402	17-11-92
		US-A- 5266569	30-11-93
		US-A- 5391763	21-02-95
		AT-T- 124040	15-07-95
		CA-A- 2023217	17-02-91
		CA-A- 2127561	17-02-91
		CN-A,B 1049501	27-02-91
		CZ-A- 9004027	17-04-96
		DE-D- 69020262	27-07-95
		DE-T- 69020262	26-10-95
		EG-A- 19251	29-09-94
		EP-A- 0413455	20-02-91
		ES-T- 2074131	01-09-95
		IL-A- 95331	31-07-95
		JP-A- 7149758	13-06-95
		JP-B- 8019099	28-02-96
		JP-A- 3086875	11-04-91
		JP-B- 7002734	18-01-95
		PL-B- 166381	31-05-95
		RU-C- 2049777	10-12-95
-----			
US-A-5164402	17-11-92	WO-A- 9102526	07-03-91
		US-A- 5266569	30-11-93
		US-A- 5391763	21-02-95
		US-A- 5229396	20-07-93
		AT-T- 124040	15-07-95
		CA-A- 2023217	17-02-91
		CA-A- 2127561	17-02-91
		CN-A,B 1049501	27-02-91
		CZ-A- 9004027	17-04-96
		DE-D- 69020262	27-07-95
		DE-T- 69020262	26-10-95
		EG-A- 19251	29-09-94
		EP-A- 0413455	20-02-91
		ES-T- 2074131	01-09-95
		IL-A- 95331	31-07-95
		JP-A- 7149758	13-06-95
		JP-B- 8019099	28-02-96
		JP-A- 3086875	11-04-91

## INTERNATIONAL SEARCH REPORT

### Information on patent family members

In                      Additional Application No                     

**PCT/IB 96/00756**

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US-A-5164402		JP-B- 7002734	18-01-95
		PL-B- 166381	31-05-95
		RU-C- 2049777	10-12-95
-----			